

Case report

Intrathoracic drainage of a perforated prepyloric gastric ulcer with a type II paraesophageal hernia

Robert A Pol*¹, Hiske W Wiersma², Bas JGL Zonneveld² and Marinus Eeftinck Schattenkerk¹

Address: ¹Department of surgery, Deventer hospital, PO Box 5001, 7416 SE, Deventer, The Netherlands and ²Department of radiology, Deventer hospital, PO Box 5001, 7416 SE, Deventer, The Netherlands

Email: Robert A Pol* - r.pol@dz.nl; Hiske W Wiersma - h.wiersma@dz.nl; Bas JGL Zonneveld - b.zonneveld@dz.nl; Marinus Eeftinck Schattenkerk - m.eeftinckschattenkerk@dz.nl

* Corresponding author

Published: 8 December 2008

Received: 16 October 2008

World Journal of Emergency Surgery 2008, **3**:34 doi:10.1186/1749-7922-3-34

Accepted: 8 December 2008

This article is available from: <http://www.wjes.org/content/3/1/34>

© 2008 Pol et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: With an incidence of less than 5%, type II paraesophageal hernias are one of the less common types of hiatal hernias. We report a case of a perforated prepyloric gastric ulcer which, due to a type II hiatus hernia, drained into the mediastinum.

Case presentation: A 61-year old Caucasian man presented with acute abdominal pain. On a conventional x-ray of the chest a large mediastinal air-fluid collection and free intra-abdominal air was seen. Additional computed tomography revealed a large intra-thoracic air-fluid collection with a type II paraesophageal hernia. An emergency upper midline laparotomy was performed and a perforated pre-pyloric gastric ulcer was treated with an omental patch repair. The patient fully recovered after 10 days and continues to do well.

Conclusion: Type II paraesophageal hernia is an uncommon diagnosis. The main risk is gastric volvulus and possible gastric torsion. Intrathoracic perforation of gastric ulcers due to a type II hiatus hernia is extremely rare and can be a diagnostic and treatment challenge.

Case report

A 61-year old Caucasian man with a previous medical history of a peptic ulcer, which was treated with a proton pump inhibitor (PPI) and H. pylori eradication, presented with acute abdominal pain after several days of extensive vomiting and abdominal discomfort. On physical examination we saw a sick man with a grey appearance. His blood pressure was 190/120 mm Hg with a pulse rate of 100 beats/minute. Upon examination the abdomen was diffusely painful with guarding and rebound tenderness. Laboratory studies showed a normal renal function and electrolytes, a normal white blood cell count of $7,2 \times 10^3$

($4,0-10,0$) and a slightly elevated c-reactive protein level of 25 mg/l (-10). On a conventional x-ray of the chest (Figure 1) a large mediastinal air-fluid collection and the suspicion of free intra-abdominal air, suggestive of a hiatal hernia and intra-abdominal perforation, was seen. In the differential diagnosis a Boerhaave syndrome was suspected. Additional computed tomography (CT) (Figure 2 and 3) revealed the aforementioned intra-abdominal free air and a large intra-thoracic air-fluid collection with a type II paraesophageal hernia. An emergency upper midline laparotomy was performed because of the septic profile and the suspicion of a perforated gastric ulcer. This




Figure 1

Chest x-ray at the emergency department. Posterior-anterior conventional radiograph of the chest with an intrathoracic air-fluid collection. Air-bubble in stomach. Free-intraperitoneal air inferior of both hemidiaphragms. Image suspect of stomach/bowel perforation and partial intrathoracic positioned stomach.

revealed a perforated pre-pyloric gastric ulcer, which was treated with an omental patch repair. The greater omentum and fundus of the stomach were partially located in the paraesophageal hernia (analogous to the CT findings) and repositioned intraabdominally. The intrathoracic collection was drained and extensively cleaned. The hiatal hernia was identified but left untreated, at the surgeon's

discretion, because of the anticipated high risk of infection and abscess formation. Postoperatively broad-spectrum antibiotics (metronidazol/cefazoline) were continued for five days and a high dose of a proton-pump inhibitor was started. The patient fully recovered after 10 days and continues to do well.




Figure 2

CT-scan. Coronal reconstruction CT-slice (3.7 mm). Intrathoracic mass consisting of mesenterial fat, free-intraperitoneal fluid and free-intraperitoneal air (1). On the left-side of this mass the esophagus is seen with a nasogastric tube (2) indicating a right-sided para-esophageal hernia with free intra peritoneal air and fluid. Intra-abdominal positioned stomach (3).




Figure 3

CT-scan. Axial 5 mm CT-slice after i.v. contrast admission. This slice shows an intra-abdominal situated stomach with nasogastric tube (1); esophagus with NGT (2); right-sided para-esophageal hernia with intraperitoneal fat (a), free-fluid (b) and free-air (c) (3); intra-peritoneal free-air (4).

Discussion

Type II paraesophageal hernias are an uncommon diagnosis and occur in less than 5% of all hiatal hernias [1]. The etiology is still unclear but previous surgical interventions, such as antireflux procedures or partial gastrectomies, have been recognized as a known risk factor. Due to progressive enlargement of the phrenoesophageal membrane, the greater curvature of the stomach tends to roll up into the thorax. Eventually, the whole stomach herniates, forming an upside-down intrathoracic stomach [2].

Most patients with a type II hernia are asymptomatic or have mild gastroesophageal reflux disease (GERD) and are diagnosed during upper gastrointestinal endoscopy.

The most important complications are gastric volvulus or bleeding from gastric ulcerations or erosions (Cameron lesions) [2,3]. The gold standard for gastric volvulus is open laparotomy with detorsion and anterior gastropexy, with or without a Nissen fundoplication [4].

Cameron lesions are linear gastric ulcers or erosions on the mucosal folds at the diaphragmatic impression in patients with a large hiatal herniam [3,5]. Unlike the prepyloric ulcer in this case, Cameron ulcers are located on the lesser curvature of the stomach. Treatment is primarily medical with acid suppressants and prokinetic agents [2,5].

To our knowledge only 5 case reports have been published reporting perforated gastric ulcers in combination with a paraesophageal hernia and just 2 cases reporting a perforated duodenal ulcer [6-12]. Normally, once a paraesophageal hernia is identified, it should be treated surgically with reduction of the herniated stomach with gastropexy to prevent reherniation and herniorraphy (or prosthetic mesh) of the diaphragm [13,14]. Debate exists whether or not an antireflux procedure is necessary. In this case no herniorraphy or mesh repair was carried out due to the anticipated high risk of infection and abscess formation. In the author's opinion risk of complications due to additional procedures should be avoided in view of the known high mortality of a perforated gastric ulcer in combination with a type II hiatus hernia.

Consent

Written informed consent was obtained from the patient for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interest

The authors declare that they have no competing interests.

Authors' contributions

RP drafted the manuscript as well as to the analysis of the literature. HW contributed to the conception and design of the manuscript, interpreted the radiology results, provided the figures and legends and revised the final manuscript. BZ and MES revised the final manuscript and provided important suggestions. All authors read and approved the final manuscript

References

1. Predicts G, Hinder RA: **Paraesophageal hiatal hernia.** Hernia, Nyhus LM, Condon RE, JB Lippincott, Philadelphia; 1995:544.
2. Weston AP: **Hiatal hernia with Cameron ulcers and erosions.** *Gastrointest Endosc Clin N Am* 1996, **6**:671.
3. Maganty K, Smith RL: **Cameron lesions; Unusual cause of gastrointestinal bleeding and anemia.** *Digestion* 2008, **77**:214-217.
4. Chau B, Dufel S: **Gastric volvulus.** *Emer Med J* 2007, **24**:446-447.
5. Cameron AJ, Higgins JA: **Linear gastric erosions. A lesion associated with large diaphragmatic hernia and chronic blood loss anemia.** *Gastroenterology* 1986, **91**:338-342.
6. Maruyama T, Fukue M, Imamura F, Nozue M: **Incarcerated paraesophageal hernia associated with perforation of the fundus of the stomach: report of a case.** *Surg Today* 2001, **31**:454-457.
7. Rakic S, Hissink RJ, Schiff BW: **Perforation of gastric ulcer associated with paraesophageal hernia causing diffuse peritonitis.** *Dig Surg* 2000, **17**:83-84.
8. Zarowitz H, Grayzel DM: **Paraesophageal hiatus hernia with gastric hemorrhage and perforation.** *Gastroenterology* 1950, **14**:314-316.
9. Kleitsch WP: **Catastrophic complications of hiatus hernia.** *AMA Arch Surg* 1952, **65**:665-672.
10. Hirshberg A, Schein M: **Gastric ulcer in diaphragmatic hernia.** *Surg Gynecol Obstet* 1992, **174**:165-171.
11. Ekelund M, Ribbe E, Willner J, Zilling T: **Perforated peptic duodenal ulcer in a paraesophageal hernia – a case report of a rare surgical emergency.** *BMC surgery* 2006, **6**:1.
12. Hanif Shiwani M, Thornton M: **Supradiaphragmatic perforated duodenal ulcer in a giant hiatus hernia.** *Can J Surg* 2008, **51**(4):E89-E90.
13. Hill LD: **Incarcerated paraesophageal hernia. A surgical emergency.** *Am J Surg* 1973, **126**(2):286-291.
14. Menguy R: **Surgical management of large paraesophageal hernia with complete intrathoracic stomach.** *World J Surg* 1988, **12**:415.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

